

PRODUCT INFORMATION

MboII

#ER0822 1500 U**Lot:** _____ **Expiry Date:** _____5'... **G A A G A (N) 8**↓...3'3'... **C T T C T (N) 7**↑...5'Concentration: 5 U/ μ LSource: *E.coli* that carries the cloned *mboII*R gene from *Moraxella bovis*Supplied with: 1 mL of 10X Buffer B
1 mL of 10X Buffer Tango**Store at -20°C**

BSA included

www.thermoscientific.com/onebio

RECOMMENDATIONS

1X Buffer B (for 100% MboII digestion)10 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 0.1 mg/mL BSA.**Incubation temperature**

37°C.

Unit DefinitionOne unit is defined as the amount of MboII required to digest 1 μ g of lambda DNA *dam*⁻ in 1 hour at 37°C in 50 μ L of recommended reaction buffer.**Dilution**

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

Double DigestsThermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of TangoTM Buffer. Please refer to www.thermoscientific.com/doubledigest to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/mL BSA.

Storage Buffer

Mboll is supplied in: 10 mM Tris-HCl (pH 7.5 at 25°C), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.2 mg/mL BSA and 50% glycerol.

Recommended Protocol for Digestion

- Add:

nuclease-free water	16 µL
10X Buffer B	2 µL
DNA (0.5-1 µg/µL)	1 µL
Mboll	0.5-2 µL*
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours*.

The digestion reaction may be scaled either up or down.

Recommended Protocol for Digestion of PCR Products

Directly after Amplification

- Add:

PCR reaction mixture	10 µL (~0.1-0.5 µg of DNA)
nuclease-free water	18 µL
10X Buffer B	2 µL
Mboll	1-2 µL*
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours*.

* See Star Activity.

Thermal Inactivation

Mboll is inactivated by incubation at 65°C for 20 min.

ENZYME PROPERTIES

Enzyme Activity in Thermo Scientific REase Buffers, %

B	G	O	R	Tango	2X Tango
100	50-100	20-50	0-20	50-100	20-50

Star Activity

An excess of Mboll (15 U/µg DNA x 1 hour) may result in star activity.

Methylation Effects on Digestion

Dam: may overlap – blocked.

Dcm: never overlaps – no effect.

CpG: may overlap – no effect.

EcoKI: never overlaps – no effect.

EcoBI: may overlap – no effect.

Stability during Prolonged Incubation

A minimum of 1 unit of enzyme is required for complete digestion of 1 µg of lambda DNA in 16 hours at 37°C.

Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
130	11	11	8	7/8	9	11

For **CERTIFICATE OF ANALYSIS** see back page

Note

- Mboll is blocked by overlapping *dam* methylation. To avoid *dam* methylation, use a *dam*⁻, *dcm*⁻ strain such as GM2163 (#M0099).
- Mboll produces DNA fragments that have a single-base 3'-extension which are more difficult to ligate than blunt-ended fragments.
- Mboll may remain associated with the cleaved DNA. This may cause DNA band shifting during electrophoresis. To avoid atypical DNA band patterns, use the 6X DNA Loading Dye&SDS Solution (#R1151) for sample preparation or heat the digested DNA in the presence of SDS prior to electrophoresis.

CERTIFICATE OF ANALYSIS

Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 10-fold overdigestion with Mboll (10 U/μg lambda DNA *dam*⁻ x 1 hour) (see Star Activity).


Ligation and Recleavage (L/R) Assay

The ligation and reclavage assay was replaced with LO test after validating experiments showed LO test ability to trace nuclease and phosphatase activities with sensitivity that is higher than L/R by a factor of 100.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of Mboll for 4 hours.

Quality authorized by:

 Jurgita Zilinskiene

PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.thermoscientific.com/onebio for Material Safety Data Sheet of the product.

© 2012 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.

